



## UWS Academic Portal

### The effects of microplastic on freshwater *Hydra attenuata* morphology & feeding

Murphy, Fionn; Prades, Laura; Ewins, Ciaran; Quinn, Brian

Published: 22/05/2016

[Link to publication on the UWS Academic Portal](#)

#### *Citation for published version (APA):*

Murphy, F., Prades, L., Ewins, C., & Quinn, B. (2016). *The effects of microplastic on freshwater Hydra attenuata morphology & feeding*. Poster session presented at SETAC Europe 26th Annual Meeting, Nantes, France.

#### **General rights**

Copyright and moral rights for the publications made accessible in the UWS Academic Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

#### **Take down policy**

If you believe that this document breaches copyright please contact [pure@uws.ac.uk](mailto:pure@uws.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



# The effects of microplastic on freshwater *Hydra attenuata* feeding & morphology

Fionn Murphy<sup>1</sup>, Laura Prades<sup>1</sup>, Ciaran Ewins<sup>1</sup>, & Brian Quinn<sup>1</sup>

<sup>1</sup>Institute of Biomedical & Environmental Health Research (IBEHR), University of the West of Scotland, Paisley, Scotland

E-mail contact: [fionn.g.murphy@gmail.com](mailto:fionn.g.murphy@gmail.com)




## Introduction

The issue of microplastic (MP) pollution has been primarily focused on the marine environment with comparatively little attention given to freshwater (Eerkes-Medrano et al., 2015). *Hydra attenuata* is a freshwater cnidarian inhabiting slow moving permeant waterbodies and play an important role in regulating the planktonic structure of these habitats (Burnett, 1973). Hydra have been used widely in bioassays to test the toxicity of numerous environmental contaminants (Quinn et al., 2008). There are currently no standardised tests for assessing the impact of microplastics. Here we investigate the potential impact MP have on the feeding and morphology of *Hydra* towards the development of a standardised bioassay.

## Materials & Methods

- 18 *Hydra* placed in 0.5 ml Eppendorf tubes
- Exposed to MP extracted from face wash
- Concentrations: Control, 0.01, 0.02, 0.04, 0.08 g/ml
- 10 *Artemia* were added to determine feeding rates
- Ingestion of MP was also recorded
- Morphology scored 10-1 (Wilby, 1988) after 3, 24, 48 & 96 hrs
- Score 8-6 reversible signs of intoxication
- 5-0 irreversible, leads to disintegration of the *Hydra*



## References

BURNETT, A. L. 1973. *Biology of Hydra*, Academic Press.

EERKES-MEDRANO, D., THOMPSON, R. C. & ALDRIDGE, D. C. 2015. Microplastics in freshwater systems: A review of the emerging threats, identification of knowledge gaps and prioritisation of research needs. *Water Research*, 75, 63-82.

QUINN, BRIAN, Francois Gagné, and Christian Blaise. "Validation of a multi-well plate toxicity test to assess feeding behaviour of the cnidarian, *Hydra attenuata*." *Fresenius Environmental Bulletin* 16.9 (2007): 1100-1107.

Wilby, O. K. "The Hydra regeneration assay." *Proceedings of workshop organised by Association Francaise de Teratologie*. 1988.


## Results

**Feeding 30 min:** Mean No. of *Artemia* ingested was significantly lower in 0.02 & 0.08 g/ml compared to the Ctrl.

**Feeding 60 min:** Mean No. of *Artemia* ingested was significantly lower in 0.04 & 0.08 g/ml compared to the Ctrl.

**MP Ingestion:** Increased as the conc. of MP increased, ranged from 0 to 10 pieces of MP. High ingestion rates also caused hydra to float.

**Morphology:** Non lethal changes observed to the morphology after 3 hrs.



## Conclusion

- MP are ingested by *Hydra attenuata*
- The presence of MP significantly reduces the feeding rate of *Hydra attenuata*
- Feeding is an ecologically relevant endpoint
- MP is ubiquitous in the environment
- What are the effects of other polymers and types (fibres, films....etc.)?

## Acknowledgements

The authors thank St-Lawrence Centre (SCL) of Montreal, Quebec for providing the *Hydra* used in this study and SETAC for providing a travel grant to attend SETAC Europe 2016